

Working with tidy data in R: tidyverse

Fundamental actions on data tables:

- choose rows — `filter()`
- choose columns — `select()`
- make new columns — `mutate()`
- arrange rows — `arrange()`
- calculate summary statistics — `summarize()`
- work on groups of data — `group_by()`

We can combine these verbs using the pipe operator: %>%

Standard R:

```
> mean(iris$Sepal.Length)
[1] 5.843333
```

With pipe:

```
> iris$Sepal.Length %>% mean()
[1] 5.843333
```

We can combine these verbs using the pipe operator: %>%

Standard R:

```
> head(iris)
```

	Sepal.Length	Sepal.Width	Petal.Length	Petal.Width	Species
1	5.1	3.5	1.4	0.2	setosa
2	4.9	3.0	1.4	0.2	setosa
3	4.7	3.2	1.3	0.2	setosa
4	4.6	3.1	1.5	0.2	setosa
5	5.0	3.6	1.4	0.2	setosa
6	5.4	3.9	1.7	0.4	setosa

We can combine these verbs using the pipe operator: %>%

With pipe:

```
> iris %>% head()
```

	Sepal.Length	Sepal.Width	Petal.Length	Petal.Width	Species
1	5.1	3.5	1.4	0.2	setosa
2	4.9	3.0	1.4	0.2	setosa
3	4.7	3.2	1.3	0.2	setosa
4	4.6	3.1	1.5	0.2	setosa
5	5.0	3.6	1.4	0.2	setosa
6	5.4	3.9	1.7	0.4	setosa

Combining pipe and assignment

These two lines do the same thing:

```
> mean_length <- mean(iris$Sepal.Length)
> mean_length <- iris$Sepal.Length %>% mean()
> mean_length
[1] 5.843333
```

Pipe example 1: count how many herbivores of different orders there are in `msleep`

Pipe example 1: count how many herbivores of different orders there are in `msleep`

```
msleep %>%  
  filter(vore == "herbi")
```

Pipe example 1: count how many herbivores of different orders there are in `msleep`

```
msleep %>%  
  filter(vore == "herbi") %>%  
  group_by(order)
```


Pipe example 1: count how many herbivores of different orders there are in `msleep`

```
msleep %>%  
  filter(vore == "herbi") %>%  
  group_by(order) %>%  
  summarize(count = n())
```

Pipe example 1: count how many herbivores of different orders there are in `msleep`

```
msleep %>%  
  filter(vore == "herbi") %>%  
  group_by(order) %>%  
  summarize(count = n()) %>%  
  arrange(desc(count))
```

Pipe example 1: count how many herbivores of different orders there are in `msleep`

```
msleep %>%  
  filter(vore == "herbi") %>%  
  group_by(order) %>%  
  summarize(count = n()) %>%  
  arrange(desc(count))
```

	order	count
1	Rodentia	16
2	Artiodactyla	5
3	Perissodactyla	3
4	Hyracoidea	2
5	Proboscidea	2
6	Diprotodontia	1
7	Lagomorpha	1
8	Pilosa	1
9	Primates	1

Pipe example 2: What is total day time for each animal in `msleep`?

Pipe example 2: What is total day time for each animal in `msleep`?

```
msleep %>%  
  mutate(total_day_time = awake + sleep_total)
```

Pipe example 2: What is total day time for each animal in `msleep`?

```
msleep %>%  
  mutate(total_day_time = awake + sleep_total) %>%  
  select(name, total_day_time)
```

Pipe example 2: What is total day time for each animal in `msleep`?

```
msleep %>%  
  mutate(total_day_time = awake + sleep_total) %>%  
  select(name, total_day_time)
```

	name	total_day_time
1	Cheetah	24.00
2	Owl monkey	24.00
3	Mountain beaver	24.00
4	Greater short-tailed shrew	24.00
5	Cow	24.00
6	Three-toed sloth	24.00
7	Northern fur seal	24.00
8	Vesper mouse	24.00
9	Dog	24.00
10	Roe deer	24.00

Pipe example 3: What is the median awake time of different orders in `msleep`?

Pipe example 3: What is the median awake time of different orders in `msleep`?

```
msleep %>%  
  group_by(order)
```

Pipe example 3: What is the median awake time of different orders in `msleep`?

```
msleep %>%  
  group_by(order) %>%  
  summarize(med_awake = median(awake))
```

Pipe example 3: What is the median awake time of different orders in `msleep`?

```
msleep %>%  
  group_by(order) %>%  
  summarize(med_awake = median(awake)) %>%  
  arrange(med_awake)
```

Pipe example 3: What is the median awake time of different orders in `msleep`?

```
msleep %>%  
  group_by(order) %>%  
  summarize(med_awake = median(awake)) %>%  
  arrange(med_awake)
```

	order	med_awake
1	Chiroptera	4.20
2	Didelphimorphia	5.30
3	Cingulata	6.25
4	Afrosoricida	8.40
5	Pilosa	9.60
6	Rodentia	11.10
7	Diprotodontia	11.60
8	Soricomorpha	13.70
9	Carnivora	13.75
10	Erinaceomorpha	13.80